BLOOD SCIENCES DEPARTMENT OF CLINICAL BIOCHEMISTRY



Title of Document: BNP Fact Sheet

Q Pulse Reference Nº: BS/CB/DCB/GEN/10

Authoriser: Peter Beresford

Version No: 2

BNP fact sheet

The BNP (NTproBNP) assay was introduced at North Bristol NHS Trust to support the BNSSG Suspected Heart Failure Pathway. The following will help you understand the test and when and how to request appropriately.

What is NTproBNP

Brain Natriuretic Peptide (BNP) is a hormone released by the heart when the ventricles are stretched e.g. by fluid overload. The hormone then causes fluid and sodium loss in the urine and mild vasodilation. In heart failure BNP levels are raised, and increase in level according to New York Heart Association classification. If BNP is normal it generally rules out heart failure. (Negative Predictive Value 97%)

When should I request a BNP test?

The NICE guideline CG108 recommends that NTproBNP is measured in patients with suspected heart failure. BNP testing will not be indicated however when there is:

A previous history of MI

A confirmed diagnosis of heart failure

Bradycardia or tachycardia (<60 or >100bpm)

Atrial fibrillation

Significant abnormality on ECG

Note: A history of previous MI and symptoms suggestive of heart failure should prompt urgent referral for Echo without delay.

Is there anything else that can affect BNP levels?

Left ventricular hypertrophy, right ventricular overload, ischaemia, tachycardia, hypoxaemia, PE, sepsis, COPD, diabetes, liver cirrhosis, age >70 and eGFR <60 ml/min can all increase BNP.

Obesity, diuretics, ACE inhibitors, beta blockers, angiotensin receptor antagonists and aldosterone antagonists can lead to falsely low levels although it is felt unlikely to impact on screening but should be factored into clinical judgement when interpreting test results.

How do I do the test?

NT proBNP is analysed by Clinical Biochemistry at Southmead Hospital. The sample should be collected into a gold top SST tube and sent to the laboratory on the day of collection. Requests should be made electronically via the ICE system where available or by paper request.

What to do with the results

BLOOD SCIENCES DEPARTMENT OF CLINICAL BIOCHEMISTRY



Version N^o: 2

Title of Document: BNP Fact Sheet

Q Pulse Reference N°: BS/CB/DCB/GEN/10


Authoriser: Peter Beresford

The BNSSG Suspected Heart Failure pathway recommends the cut-offs and care pathway

below.

Heart failure is unlikely if the NTproBNP level is:

<400 pg/mL



NTproBNP <400 pg/mL

Heart Failure unlikely

Consider other diagnosis

NTproBNP **400 – 2000 pg/mL**

PRIMARY CARE to request ECG at their *local* secondary care hospital

NTproBNP >2000 pg/mL

Refer to Sirona HF service for a review at the one stop Echo/Consultant clinic within 2 weeks

DO NOT REQUEST ECHO

Page 2 of 2